

國立中央大學98學年度碩士班考試入學試題卷

所別：工業管理研究所碩士班 甲組

科目：統計學 共 3 頁 第 1 頁

乙組

*請在試卷答案卷（卡）內作答

1. An automotive component has been designed to withstand certain stressed. It is known from the past experience that, because of variation in loading, the stress on the component is normally distributed with a mean of 30000 kPa and a standard deviation of 3000 kPa. The strength of the component is also random because of variation in the material characteristics and the dimensional tolerances. It has been found that the strength is normally distributed with a mean of 40000 kPa and a standard deviation of 4000 kPa. Determine the reliability of the component. (15%)
2. In humans there is a blood group, the MN group, that is composed of individuals having one of the three blood types M, MN, and N. Type is determined by two alleles, and there is no dominance, so the three possible genotypes give rise to three phenotypes. A population consisting of individuals in the MN group is in equilibrium if $P(M)=p_1=\theta^2$ $P(MN)=p_2=2\theta(1-\theta)$ $P(N)=p_3=(1-\theta)^2$ for some θ . Suppose a sample from such population yields the results shown as

Type	M	MN	N	
Observed	125	225	150	N=500

- (a) Find the MLE for θ (10%)
- (b) Do the data provide the sufficient evidence that the population is in equilibrium status? $\alpha=0.05$. (10%)
3. Nine weeks were randomly selected and the absentee rate (percentage of workers absent) determined for each day (Monday through Friday) of the workweek. The data are reproduced in the table. Conduct a test to compare the distributions of absentee rates for the 5 days of the workweek at $\alpha=0.05$. (15%)

Week	Monday	Tuesday	Wednesday	Thursday	Friday
1	5.3	0.6	1.9	1.3	1.6
2	12.9	9.4	2.6	0.4	0.5
3	0.8	0.8	5.7	0.4	1.4
4	2.6	0.0	4.5	10.2	4.5
5	23.5	9.6	11.3	13.6	14.1
6	9.1	4.5	7.5	2.1	9.3
7	11.1	4.2	4.1	4.2	4.1
8	9.5	7.1	4.5	9.1	12.9
9	4.8	5.2	10.0	6.9	9.0

參考用

注意：背面有試題

國立中央大學98學年度碩士班考試入學試題卷

所別：工業管理研究所碩士班 甲組 科目：統計學 共 3 頁 第 2 頁
乙組 *請在試卷答案卷(卡)內作答

4. For each event E of the sample space Ω , define probability, $P(E)$, to satisfy the following three axioms:

$$(1) 0 \leq P(E) \leq 1$$

$$(2) P(\Omega) = 1$$

- (3) for any sequence of events E_1, E_2, \dots that are mutually exclusive, that is, events for which $E_i \cap E_j = \emptyset$ when $i \neq j$,

$$P\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} P(E_i)$$

Show that $P(E \cup F) = P(E) + P(F) - P(E \cap F)$, and $P\left(\bigcup_{i=1}^{\infty} E_i\right) \leq \sum_{i=1}^{\infty} P(E_i)$. (10%)

5. By the method of least squares, fit the cubic $y = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3$ to the 10 observations of (x, y) : (10%)

$$(0, 0), (0, 1), (0, 2), (-1, -2), (-1, -3), \\ (1, 1), (1, 3), (-2, -8), (2, 9), (2, 10).$$

6. Let Y be the sum of n observations of a random sample from a Poisson distribution with mean θ . Let the prior p.d.f. of θ be a gamma distribution with parameters α and β .

- (a). Given $Y = y$, find the posterior p.d.f. of θ . (10%)

- (b). If the loss function is $[\hat{\theta} - \theta]^2$, find the Bayesian point estimate $\hat{\theta}$. (10%)

- (c). Show that this $\hat{\theta}$ is a weighted average of the maximum likelihood estimate

$\frac{y}{n}$ and the prior mean $\alpha\beta$, with respective weights of $\frac{n}{n + \frac{1}{\beta}}$ and

$$\frac{\frac{1}{\beta}}{n + \frac{1}{\beta}}. (10%)$$

注意：背面有試題


國立中央大學98學年度碩士班考試入學試題卷

所別：工業管理研究所碩士班 甲組
乙組

科目：統計學 共 3 頁

*請在試卷答案卷（卡）內作答


STANDARD NORMAL DISTRIBUTION



Entries in the table give the area under the curve between the mean and z standard deviations above the mean. For example, for $z = 1.25$ the area under the curve between the mean and z is .3944.

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2518	.2549
.7	.2580	.2612	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974	.4975
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4980	.4981	.4982
2.9	.4981	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986	.4987
3.0	.4986	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990	.4990

The Chi-Square Distribution



This table is abridged and adapted from Table III in Biometrika Tables for Statisticians, edited by E. S. Pearson and H. O. Hartley. It is published here with the kind permission of the Biometrika Trustees.